Motion of test particles in a magnetized conformastatic background

Antonio C. Gutiérrez-Piñeres

Facultad de Ciencias Básicas, Universidad Tecnológica de Bolívar, Cartagena 13001, Colombia
Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, AP 70543, México, DF 04510, México
E-mail: gutierrezpac@gmail.com

Abraão J. S. Capistrano

Federal University of Latin-American Integration, Casimiro Montenegro Filho Astronomical Pole, Technological Park of Itaipu Technological Park of Itaipu, PO box 2123, Foz do Iguaçu-PR 85867-670, Brazil
E-mail: abraao.capistrano@unila.edu.br

arXiv:1510.05400v1 [gr-qc] 19 Oct 2015

A class of exact conformastatic solutions of the Einstein-Maxwell field equations is presented in which the gravitational and electromagnetic potentials are completely determined by a harmonic function only. The motion of test particles is investigated in the background of a space-time characterized by this class of solutions. We focus on the study of circular stable and unstable orbits obtained by taking account particular harmonic functions defining the gravitational potential. We show that is possible to have repulsive force generated by the charge distribution of the source. As the space-time here considered is singularity free we conclude that this phenomena is not exclusive to the case of naked singularities. Additionally, we obtain an expression for the perihelion advance of the test particles in a general magnetized conformastatic space-time.

Keywords: Exact Solutions; Circular Orbits; Perihelion Advance; Repulsive Force; Naked Singularity.

1. A conformastic solution of the Einstein-Maxwell equations

In Einstein-Maxwell gravity theory the background of a symmetric body can be described by the conformastatic metric in cylindrical coordinates [1]

$$dS^2 = -c^2 e^{2\phi} dt^2 + e^{-2\phi}(dr^2 + dz^2 + r^2 d\varphi^2), \tag{1}$$

where c is the speed of the light in the vacuum and the metric potential ϕ depends only on the r and z variables. The governing equations field are the Einstein-Maxwell equations which, for the metric (1), can be equivalently written as [a],

$$\nabla^2\phi = \nabla\phi\cdot\nabla\phi, \quad \phi_{,r}^2 = \frac{G}{c^4 r^2} e^{2\phi} A_{\varphi,z}^2, \quad \phi_{,z}^2 = \frac{G}{c^4 r^2} e^{2\phi} A_{\varphi,r}^2, \tag{2}$$

$$\phi_{,r}\phi_{,z} = -\frac{G}{c^4 r^2} e^{2\phi} A_{\varphi,r} A_{\varphi,z}, \quad \nabla\cdot(r^{-2} e^{2\phi}\nabla A_\varphi) = 0, \tag{3}$$

where the operator ∇ denotes the usual gradient in cylindrical coordinates. By supposing a solution of $\nabla^2\phi = \nabla\phi\cdot\nabla\phi$ in the functional form $\phi = \phi[U(r,z)]$, where

[a] Along this work we use the CGS units such that $k_0 \equiv 8\pi G c^{-4} = 2,07 \times 10^{-48}\ \mathrm{s^2 cm^{-1} g^{-1}}$, $G = 6.674 \times 10^{-8}\mathrm{cm^3 g^{-1} s^{-2}}$ and $c = 2.998 \times 10^{10}\,\mathrm{cm\,s^{-1}}$.

$U(r,z)$ is an arbitrary harmonic function (See Refs. 1 and 2 for more details) it is not difficult to prove that

$$\phi = -\ln(1-U), \quad A_\varphi(r,z) = \frac{c^2}{G^{1/2}}\int_0^r \tilde{r}U(\tilde{r},z)d\tilde{r}, \tag{4}$$

is a solution of the previous equations system. The electromagnetic field is of a magnetic type. In fact, can be demonstrated that the electromagnetic invariant $\mathcal{F} \equiv F^{\alpha\beta}F_{\alpha\beta} \geq 0$.The nonzero components of the electromagnetic fields read $B_i = c^2G^{-1/2}rU_{,i}$, $(i = r, z)$ where, as we know, U is an arbitrary harmonic function.

The motion of a test particle of mass m and charge q moving in a conformastatic space-time given by the line element (1) is described by the Lagrangian $\mathcal{L} = \frac{1}{2}mg_{\alpha\beta}\dot{x}^\alpha\dot{x}^\beta + \frac{q}{c}A_\alpha\dot{x}^\alpha$, where A_α is the four-potential and the dot represents differentiation with respect to the proper time. Since the Lagrangian does not depend explicitly of the variables t and φ, the following two conserved quantities exist

$$p_t = -mce^{2\phi}\dot{t} \equiv -\frac{E}{c}, \quad \text{and} \quad p_\varphi = mr^2e^{-2\phi}\dot{\varphi} + \frac{q}{c}A_\varphi \equiv L, \tag{5}$$

where E and L are, respectively, the energy and the angular momentum of the particle as measured by an observer at rest at infinity. The momentum p_α of the particle can be normalized so that $g_{\alpha\beta}\dot{x}^\alpha\dot{x}^\beta = -\Sigma$, accordingly, for the metric (1)

$$-c^2e^{2\phi}\dot{t}^2 + e^{-2\phi}(\dot{r}^2 + \dot{z}^2 + r^2\dot{\varphi}^2) = -\Sigma, \tag{6}$$

where $\Sigma = 0, c^2, -c^2$ for null, time-like, and space-like curves, respectively.

To investigate the motion of the test particle we restrict ourselves to the study of the equatorial trajectories with $z = 0$. Accordingly, by the substitution of the conserved quantities (5) into Eq.(6) we find

$$\dot{r}^2 + \Phi = \frac{E^2}{m^2c^2}, \quad \text{where} \quad \Phi(r) \equiv \frac{L^2}{m^2r^2}\left(1 - \frac{qA_\varphi}{Lc}\right)^2 e^{4\phi} + \Sigma e^{2\phi} \tag{7}$$

is an effective potential. In this work we present an analysis of the circular motion of test particles governed by the above equations. As a particular sample we consider the effective potential corresponding to a charged particle moving in the gravitational field of a punctual source placed in the coordinates origin of a magnetized conformastatic space-time. Additionally, we find an expression for the perihelion advance of the test particles in a general magnetized conformastatic space-time. This work is inspired in the Refs. 3 and 4.

2. Circular motion in the field of a punctual source in Einstein-Maxwell gravity

The motion of charged test particles is governed by the behavior of the effective potential (7). The radius of circular orbits and the corresponding values of the

energy E and angular momentum L are given by the extrema of the function Φ. Therefore, the conditions for the occurrence of circular orbits are

$$\frac{d\Phi}{dr} = 0, \quad \Phi = \frac{E^2}{m^2c^2}. \tag{8}$$

Notice that the positive value of the energy corresponds to the positive of solution $E_\pm = \pm mc\Phi^{1/2}$. An interesting particular orbit is the one in which the particle is located at rest (r_r) as seen by an observer at infinity, i.e. $L = 0$. These orbits are therefore characterized by the conditions $L = 0, \quad d\Phi/dr = 0$. For the metric (1) these conditions give us the following equation for the rest radius

$$\frac{2e^{2\phi}}{m^2c^2r^3}\left[q^2A_\varphi\left(2rA_\varphi\phi_{,r} + rA_{\varphi,r} - A_\varphi\right)e^{2\phi} + \Sigma m^2c^2r^3\phi_{,r}\right] = 0. \tag{9}$$

If $e^{2\phi} = 0$ for a orbit with a rest radius $r = r_r$ the energy of the particle is $E_r = 0$. In the case $e^{2\phi} \neq 0$ we have

$$E^{(\pm)}{}_{r\pm} = \pm mce^{\phi}\left(\Sigma + \xi^{(\pm)}{}_r\right)^{1/2}, \tag{10}$$

where

$$\xi^{(\pm)}{}_r = \frac{q^2e^{2\phi}\left[rA_{\varphi,r} \pm \left(rA_{\varphi,r} + 2A_\varphi\left(2r\phi_{,r} - 1\right)\right)\right]^2}{4m^2c^2r^2\left(2r\phi_{,r} - 1\right)^2}, \tag{11}$$

To obtain r_r we must to solve the equation (9) for r_r. The minimum radius for a stable circular orbits occurs at the inflection points of the effective potential function, thus we must solve the equation $d^2\Phi/dr^2 = 0$, using the expresion of the angular moment L of the particle in a circular orbit. From the equation (8) and the last equation we find that the radius of last stable circular orbit and the angular moment of this orbit.

The gravitational and magnetic field in the background of a source of mass M placed in the origin of coordinates in Einstein-Maxwell gravity can be obtained by consider the harmonic potential

$$U(r,z) = -\frac{GM}{c^2R}, \quad R^2 = r^2 + z^2 \tag{12}$$

in the equation (4). Accordingly, for the metric and electromagnetic potential we have

$$\phi(r,z) = -\ln\left(1 + \frac{GM}{c^2R}\right) \quad \text{and} \quad A_\varphi(r,z) = \sqrt{G}M\left(1 - \frac{z}{R}\right) \tag{13}$$

respectivly. By calculating the asymptotic behavior of the metric potential g_{tt}, the Kretschmann scalar and the electromagnetic invariant $\mathcal{F}$ we conclude that the gravitational field is asymptotically like Schwarzschild and singularity free. We notice that when we have the solution Eq.(12), we recover the extreme Reissner-Nordström solution in its standard form by writing the metric in spherical polar coordinates and replacing R by $R - GM/c^2$.

Consider the case of a charged particle moving in the fields given by (13). This means that we are considering the motion described by the following effective potential

$$\Phi(r) = \frac{c^6 r^2 (Lc - q\sqrt{G}M)^2}{m^2(c^2 r + GM)^4} + \frac{\Sigma c^4 r^2}{(c^2 r + GM)^2}. \tag{14}$$

From Eq.(9) for $q \neq 0$ we find the obvious rest radius $r_r = 0$. In this radius the energy of the particle is $E_r = 0$. Similarly we find the rest radii

$$\frac{r_{r\,(\pm)}}{M} = \frac{c^2q^2 - 2\Sigma GMm^2(\pm)\sqrt{c^2q^2\,(c^2q^2 - 8\Sigma Gm^2)}}{2\Sigma m^2 c^2}. \tag{15}$$

For the rest radius $r_{r\,(\pm)}$ we have the following values for the energy

$$E^{(\pm)}_{r\pm} = \pm\frac{mc^2\sqrt{2q^2\left[q^2 - 2Gm^2(\pm)\sqrt{q^2(q^2 - 8Gm)^2}\right]^3}}{\left[q^2 + \sqrt{q^2(q^2 - 8Gm)^2}\right]^2}, \tag{16}$$

Notice that for each value of r_r is possible to have a particle in rest with negative energy (E_{r-}). Also notice that, the rest radii are conditioned by the mass m and charge q of the test particle. For null curves these are not defined. For time-like curves ($\Sigma = c^2$) the rest radii for a charged particle are restricted by the discriminant $q^2 - 8m^2G$ in the formula (15). In fact, the rest radii take real positive values for all value of q and m except for those in the interval $(-\sqrt{8G}m, \sqrt{8G}m)$. Note that for the value of the charge $q^2 = 8m^2G$, we have $r_r = 3GM/c^2$. For this value of charge the energy of the charged test particle E_{r+} reaches its minimum value which is $+(3\sqrt{6}/8)mc^2$ whereas the energy E_{r-} reaches its maximum value, which is $-(3\sqrt{6}/8)mc^2$. Additionally, the angular momentum and the energy for a circular orbit with radius r_c are given by

$$L_{c\,\pm} = \frac{q\sqrt{G}M}{c} \mp \frac{(c^2 r_c + GM)m}{c^2}\sqrt{\frac{\Sigma GM}{c^2 r_c - GM}} \tag{17}$$

and

$$E_{c\,\pm} = \pm\frac{mc^4}{(c^2 r_c + GM)}\sqrt{\frac{\Sigma r_c^3}{c^2 r_c - GM}} \tag{18}$$

respectively. From Eq.(17) and (18) we conclude that in order to have a time-like circular orbit the charged particle must be placed in a radius $r_c > GM/c^2$. It is easy to note that for the radius $r_c = 3GM/c^2$ the angular momentum of the particle is $L_c^{\pm} = q\sqrt{G}M/c \mp \sqrt{2}/2GMm/c$. Notice that for this radius and the particular values of the charge $q = 2\sqrt{2G}m$ and $q = -2\sqrt{2G}m$ we have the corresponding values for the angular momentum $L_c^{\pm} = (1 \mp 1)\sqrt{2}GMm/(2c)$ and $L_c^{\pm} = (-1 \mp 1)\sqrt{2}GMm/(2c)$, respectively. Accordingly, for this radius and the charge $q = 2\sqrt{2G}m$ we have $L_c^{+} = L_r^{+} = 0$. Similarly, for this radius and the charge $q = -2\sqrt{2G}m$ we have $L_c^{-} = L_r^{-} = 0$. In these cases the value of the

energy of the particle is $E_{c\pm} = E_{r\pm} = \pm(3\sqrt{6}/8)mc^2$. Moreover, from the solution of the equation $d^2\Phi/dr^2 = 0$ for L and Eq.(17) we find that the last stable circular orbit satisfies the very simple equation $r(c^2r - 3GM) = 0$. Therefore the last stable circular orbit occurs in $r_{lsco} = 3GM/c^2$ and the angular momentum of the last stable circular orbit is given by $c^2L_{lsco\,\pm} = qc\sqrt{G}M \pm 2\sqrt{2\Sigma}GMm$ As we can see, the angular momentum and the energy are conditioned by the mass m and charge q of the test particle. As we note, for space-like curves the angular momentum of the last stable circular orbit it is not defined, whereas for a null curves the angular momentum of the last stable circular orbit is $L_{lsco\,\pm} = q\sqrt{G}M/c$. The angular momentum of a charged particle in a time-like circular stable orbit is given by $L_{lsco\,\pm} = (q\sqrt{G} \pm 2\sqrt{2}Gm)M/c$. Accordingly we can see that if the charge of the particle is $q = 2\sqrt{2G}m$ then $L_{lsco-} = L_{c+} = L_r = 0$. Analogously, if $q = -2\sqrt{2G}m$ then $L_{lsco+} = L_{c-} = L_r = 0$. Then we deduce that the particle can be placed in a last stable circular orbit of radius $r = 3GM/c^2$ for some arbitrary values of the charge. Whereas the particle will be in rest in a stable position $r = 3GM/c^2$ if the value of the charge is $q = \pm 2\sqrt{2G}m$.

The energy of the charged particle can be written in terms of the its effective potential Φ as (see Eq. (8)) $E = \pm mc\sqrt{\Phi}$, with Φ given by Eq.(14). At the infinitum, the effective potential (energy) tends to a constant. As we know, the radius for the last stable orbit is given by $c^2r = 3GM$ It is possible then, for this radius and value of the charge $q/m = 2\sqrt{2G}$, to have a particle a particle in rest or in an orbit with angular momentum $4\sqrt{2}G/c$ ("clockwise" motion). A similar result is obtained if the charge is negative. The angular momentum in such case corresponds to a charged particle with angular momentum $-4\sqrt{2}G/c$ ("counterclockwise" motion). In the classical radius $r = 3GM/c^2$ circular orbits exist with zero angular momentum. That phenomena is interpreted as a consequence of the repulsive force generated by the charge distribution. As the space-time here considered is free of singularities we conclude that this phenomena is not exclusive to the case of naked singularities.

3. The perihelion advance in a conformastatic magnetized spacetime

To study the problem of the perihelion advance of a test charged particle in a conformastatic spacetime in presence of a magnetic field we start of the equation (6). We restrict the motion of the particle to the plane $z = 0$. Then we have the expression

$$\frac{d^2u}{d\varphi^2} + u^2 = F(u), \quad u = 1/r, \tag{19}$$

where

$$F(u) \equiv \frac{1}{2}\frac{dG}{du}, \qquad G(u) \equiv \frac{1}{\left(1 - \frac{qA_\varphi}{cL}\right)^2}\left[\frac{E^2}{c^2L^2}(1-U)^4 - \frac{\Sigma m^2}{L^2}(1-U)^2\right]. \tag{20}$$

Accordingly, by following a procedure in the same way as in Ref. 5, we have for the resulting perihelion advance $\delta\varphi = \pi(dF/du)_{u=u_0}$, where u_0 is the radius of a nearly circular orbit, which is given by the roots of the equation $F(u_0) = u_0$. By inserting Eq.(12) into Eq.(20) we obtain for $F(u)$

$$F(u) = \left[\frac{2E^2GM}{c^4L^2}\left(1+\frac{GM}{c^2}u\right)^3 - \frac{\Sigma m^2 GM}{c^2L^2}\left(1+\frac{GMu}{c^2}\right)\right]\left(1-\frac{q\sqrt{G}M}{cL}\right)^{-2},$$

accordingly, we find that the perihelion advance in the orbit of the test particles in this kind of spacetime is $\delta\varphi = \pi(\psi_0 - k_2^2)/Q^2$, where

$$\psi_0 \equiv \frac{\left[6\left(Q^2+k_2^2\right) + \left[54Q^2k_1\left(-1+\sqrt{1-\frac{6\left(Q^2+k_2^2\right)^3}{81Q^4k_1^2}}\right)\right]^{2/3}\right]^2}{6\left[54Q^2k_1\left(-1+\sqrt{1-\frac{6\left(Q^2+k_2^2\right)^3}{81Q^4k_1^2}}\right)\right]^{2/3}},$$

with

$$K_1^2 = \frac{E^2G^2M^2}{c^6L^2}, \quad K_2^2 = \frac{\Sigma m^2G^2M^2}{c^4L^2}, \quad \text{and} \quad Q^2 = \left(1-\frac{q\sqrt{G}M}{cL}\right)^2.$$

Notice that when $q = 0$ we get the case in which $\delta\varphi$ describe the perihelion advance of a neutral particle. As we recover the extreme Reissner-Nordström solution in its standard form by writing the metric in spherical polar coordinates and replacing R by $R - GM/c^2$ in the metric then is posible to perform a comparison between the results presented here and the corresponding results presented in another works. In addition, along this work we have used the CGS units, in such a way that a comparison with the observational data can be applied.

References

1. A. C. Gutiérrez-Piñeres, G. A. González and H. Quevedo, Conformastatic disk-haloes in einstein-maxwell gravity, *Phys. Rev. D* **87**, p. 044010 (Feb 2013).
2. A. C. Gutiérrez-Piñeres, Conformastationary disk-haloes in einstein-maxwell gravity, *General Relativity and Gravitation* **47** (2015).
3. D. Pugliese, H. Quevedo and R. Ruffini, Motion of charged test particles in reissner-nordström spacetime, *Phys. Rev. D* **83**, p. 104052 (May 2011).
4. D. Pugliese, H. Quevedo and R. Ruffini, Circular motion of neutral test particles in reissner-nordström spacetime, *Phys. Rev. D* **83**, p. 024021 (Jan 2011).
5. T. Harko, Z. Kovács and F. S. N. Lobo, Solar system tests of hořava–lifshitz gravity, *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences* (2010).